



Research Paper

SOME INEQUALITIES OF REFORMULATED AND ENTIRE INF-SOMBOR INDEX

JALAPPA¹ MANOHAR S. H.^{1,2} AND BOREGOWDA H. S.^{1*}

¹ Department of Studies and Research in Mathematics, Tumkur University, Jnanasiri campus, Bidarakatte, Tumakuru-572118. metijalappa@gmail.com

¹ Department of Studies and Research in Mathematics, Tumkur University, Jnanasiri campus, Bidarakatte, Tumakuru-572118. bgsamarasa@gmail.com

¹ Department of Studies and Research in Mathematics, Tumkur University, Jnanasiri campus, Bidarakatte, Tumakuru-572118. manoharsh146@gmail.com

² Department of Mathematics, Government First Grade College Madhugiri-572132. manoharsh146@gmail.com

ARTICLE INFO

Article history:

Received: 26 December 2024

Accepted: 24 April 2025

Communicated by Alireza Naghipour

Keywords:

Platt number

Inf Sombor index

Reformulated Inf Sombor index

Entire Inf Sombor index.

MSC:

05C07; 05C09; 05C92.

ABSTRACT

The concept of the Sombor index was extended by Reti *et. al.* [2] by introducing a p -Sombor index which can be seen as the p norm of the vector $x = (d(u), d(v))$ as $p \rightarrow \infty$, $\|x\|_\infty = \max\{d(u), d(v)\}$. Inspired by this, we defined indices namely, Reformulated Inf- Sombor index, Entire Inf-Sombor index, and KG Inf-Sombor index. Also, we present lower and upper bounds by using some graph parameters and obtain exact values of these new topological indices in some graph families. Further, we evaluated the statistical behavior of these indices after computing index values for different types of dendrimers for various growth values k .

1. INTRODUCTION

We only take into account undirected, simple, and finite graphs. Let $\delta = \delta(G)$ and $\Delta = \Delta(G)$ be the minimum and maximum vertex degree of the graph G respectively. $g \sim h$ means that the edges g and h are adjacent, i.e., they share a common end-vertex in G and let $u \in$

*Address correspondence to H. S. Boregowda; Department of Studies and Research in Mathematics, Tumkur University, Jnanasiri campus, Bidarakatte, Tumakuru-572118, E-mail: bgsamarasa@gmail.com.

Topological indices	Mathematical expressions
Forgotten index [8, 9]	$F(G) = \sum_{uv \in E(G)} [d(u)^2 + d(v)^2]$
Randic index [11, 10]	$R(G) = \sum_{uv \in E(G)} \frac{1}{\sqrt{d(u).d(v)}}$
First Zagreb index [5]	$M_1(G) = \sum_{uv \in E(G)} [d(u) + d(v)]$
Second Zagreb index [6, 7]	$M_2(G) = \sum_{uv \in E(G)} [d(u).d(v)]$
Entire Randic index [17]	$ER(G) = \sum_{\{g,h\} \subseteq B(G)} \frac{1}{\sqrt{d(g)d(h)}}$
p-Sombor index [2]	$SO_p(G) = \sum_{uv \in E(G)} [d(u)^p + d(v)^p]^{1/p}$
Sombor index [1]	$SO(G) = \sum_{uv \in E(G)} [\sqrt{d(u)^2 + d(v)^2}]$
Inf-Sombor index [12]	$SO_\infty(G) = \sum_{uv \in E(G)} \max\{d(u), d(v)\}$
KG Sombor index [22]	$KG(G) = \sum_{ue} [\sqrt{d(u)^2 + d(e)^2}]$
Reformulated Sombor index [20]	$RS(G) = \sum_{g \sim h} \sqrt{[d(g)^2 + d(h)^2]}$
Reformulated first Zagreb index [21]	$REM_1(G) = \sum_{g \sim h} [d(g) + d(h)]$
Platt index [13]	$Pl(G) = \sum_{u \in V(G)} d(u) (d(u) - 1)$
Reformulated second Zagreb index [21]	$REM_2(G) = \sum_{g \sim h} [d(g)d(h)]$
First Entire Zagreb index [23]	$EM_1(G) = \sum_{\{g,h\} \subseteq B(G)} [d(g) + d(h)]$
Second Entire Zagreb index [23]	$EM_2(G) = \sum_{\{g,h\} \subseteq B(G)} [d(g)d(h)]$
Entire Sombor index [18, 19]	$ES(G) = \sum_{\{g,h\} \subseteq B(G)} \sqrt{[d(g)^2 + d(h)^2]}$
Reformulated Inverse sum Indeg index [29]	$ISI_e(G) = \sum_{g \sim h} \frac{d(g)d(h)}{d(g)+d(h)}$

TABLE 1. Topological indices and their mathematical expressions

$V(G)$, $e \in E(G)$ then ue represents u is incident to e . If there is a path connecting each pair of vertices in a graph G , then that graph is said to be connected. The degree of a vertex v , shown by $d(v)$, is the number of vertices that are adjacent to it. Furthermore, $d(e) = d(u) + d(v) - 2$ defines the edge ($e = uv$) degree. For further terminology and notation pertaining to graph theory, see [3, 4]. The Sombor index, defined as $\sum_{uv \in E(G)} [\sqrt{d(u)^2 + d(v)^2}]$, was initially defined by Gutman [1] in 2021. It is based on the degree radius of an edge $e = uv \in E(G)$. The concept of the Sombor index was extended by Reti *et. al.* [2]. by introducing a p -Sombor index which can be seen as the p norm of the vector $x = (d(u), d(v))$ as $p \rightarrow \infty$, $\|x\|_\infty = \max\{d(u), d(v)\}$. Inspired by this, we defined indices namely, Reformulated Inf- Sombor index, Entire Inf-Sombor index, and KG Inf-Sombor index. The reformulated inf-Sombor index is defined as $RSO_\infty(G) = \sum_{g \sim h} \max\{d(g), d(h)\}$ and Entire Inf-Sombor index which is defined as the sum of the maximum of the terms g and h , which are the two members of the set $B(G)$, where $B(G)$ is the collection of all subsets of two members $\{g, h\} \subseteq V(G) \cup E(G)$ such that g and h are adjacent or incident to each other and is represented as $ESO_\infty(G) = \sum_{\{g,h\} \subseteq B(G)} \max\{d(g), d(h)\}$. KG Sombor index is defined by Kulli *et.al.* [22] which is defined as $KG(G) = \sum_{ue} \sqrt{d(u)^2 + d(e)^2}$, motivated by this we defined an index named KG Inf-Sombor index is defined as $KG_\infty(G) = \sum_{ue} \max\{d(u), d(e)\}$.

Some well-studied topological descriptors have many applications in the field of chemical graph theory. For more details refer [25, 24, 27, 28, 34, 36, 35, 26].

2. REFORMULATED INF-SOMBOR INDEX

This section contains the precise values of the reformulated Inf-Sombor index for certain graph families.

Observation: Any isolated vertices in a graph G have no impact on $L(G)$, since $L(G)$ is defined on its edge set. It is obvious that $RSO_\infty(G) = SO_\infty(L(G))$.

Theorem 2.1. *Let G be r -regular graph. Then,*

$$RSO_\infty(G) = (r - 1)^2 nr.$$

Proof. Let G be an r -regular graph. Then,

$$\begin{aligned} RSO_\infty(G) &= \sum_{g \sim h} \max\{d(g), d(h)\} \\ &= \frac{Pl(G)}{2} 2(r - 1) \\ RSO_\infty(G) &= (r - 1)^2 nr \end{aligned}$$

□

Theorem 2.2. (1) *For complete graph K_n with $n \geq 3$, $RSO_\infty(K_n) = n(n - 1)(n - 2)^2$.*

(2) *For cycle C_n with $n \geq 3$, $RSO_\infty(C_n) = 2n$.*

(3) *For complete bipartite graph $K_{m,n}$ with $1 \leq m \leq n$, $RSO_\infty(K_{m,n}) = \frac{(m+n-2)^2 mn}{2}$.*

(4) *For path P_n with $n \geq 2$, $RSO_\infty(P_n) = 2(n - 2)$.*

(5) *For star graph S_n , with $n \geq 2$, $RSO_\infty(S_n) = \frac{(n-2)^2(n-1)}{2}$.*

2.1. Inequalities related to Reformulated Inf-Sombor index.

Theorem 2.3. *Let G be any graph. Then,*

$$Pl(G)(\delta - 1) \leq RSO_\infty(G) \leq Pl(G)(\Delta - 1).$$

Proof. Let G be a connected. Then,

$$\begin{aligned} RSO_\infty(G) &= \sum_{g \sim h} \max\{d(g), d(h)\} \\ &\leq 2(\Delta - 1) \sum_{g \sim h} 1 \\ (2.1) \quad &\leq Pl(G)(\Delta - 1). \end{aligned}$$

$$\begin{aligned} \text{Similarly, } RSO_\infty(G) &= \sum_{g \sim h} \min\{d(g), d(h)\} \\ (2.2) \quad &\geq Pl(G)(\delta - 1). \end{aligned}$$

From equations (2.1) and (2.2), we get

$$(2.3) \quad Pl(G)(\delta - 1) \leq RSO_\infty(G) \leq Pl(G)(\Delta - 1).$$

Equality holds when graph G is regular. □

Lemma 2.4. [14] *For any connected graph G with $n \geq 2$,*

(1) $2m(\delta - 1) \leq Pl(G) \leq 2m(\Delta - 1)$.

(2) $m \leq Pl(G) \leq 2m(n - 2)$.

Lemma 2.5. [15, 16] *For any graph G with $n \geq 2$, $Pl(G) = M_1(G) - 2m$.*

Corollary 2.6. *For a graph G ,*

(1) $2m(\delta - 1)^2 \leq Pl(G) \leq 2m(\Delta - 1)^2$.

$$(2) \quad m(\delta - 1) \leq RSO_\infty(G) \leq 2m(n - 2)(\Delta - 1).$$

Proof. By using Lemma 2.4 in equation (2.3), we get the results. \square

Theorem 2.7. For any graph G ,

$$\frac{1}{\sqrt{2}}RSO(G) \leq RSO_\infty(G) \leq RSO(G).$$

Proof. Let G be a graph. Then,

$$\begin{aligned} RSO(G) &= \sum_{g \sim h} \sqrt{d(g)^2 + d(h)^2} \\ &\leq \sum_{g \sim h} \sqrt{2 \max\{d(g)^2, d(h)^2\}} \\ (2.4) \quad &\leq \sqrt{2}RSO_\infty(G). \end{aligned}$$

$$\begin{aligned} \text{Consider, } \sqrt{\max\{d(g)^2, d(h)^2\}} &\leq \sqrt{d(g)^2 + d(h)^2} \\ (2.5) \quad RSO_\infty(G) &\leq RSO(G). \end{aligned}$$

From equations (2.4) and (2.5), we get $\frac{1}{\sqrt{2}}RSO(G) \leq RSO_\infty(G) \leq RSO(G)$. \square

Corollary 2.8. For any graph G ,

$$\frac{m}{2} \leq RSO_\infty(G) \leq 2m(n - 2)^2.$$

Proof. From Lemma (2.4), we have

$$\begin{aligned} (2.6) \quad \frac{Pl(G)}{2} &\leq \max\{d(e), d(f)\} \leq 2(n - 2) \cdot \frac{Pl(G)}{2} \\ \frac{m}{2} &\leq \max\{d(e), d(f)\} \leq 2m(n - 2)^2. \end{aligned}$$

\square

Corollary 2.9. For any graph G

$$\frac{M_1(G) - 2m}{2} \leq RSO_\infty(G) \leq (n - 1)(M_1(G) - 2m).$$

Proof. By using Lemma (2.5) in equation (2.6), we get the result. \square

Theorem 2.10. For any graph G ,

$$\frac{REM_1(G)}{2} \leq RSO_\infty(G) \leq REM_1(G).$$

Proof. For any integer $x, y \geq 0$

$$\begin{aligned} \frac{g + h}{2} &\leq \max\{g, h\} \leq g + h \\ \sum_{g \sim h} \frac{d(g) + d(h)}{2} &\leq \sum_{g \sim h} \max\{d(g), d(h)\} \leq \sum_{g \sim h} d(g) + d(h) \\ \frac{REM_1(G)}{2} &\leq RSO_\infty(G) \leq REM_1(G). \end{aligned}$$

\square

Theorem 2.11. Let G be any graph. Then,

$$RM_1(G) - (\Delta - 1)Pl(G) \leq RSO_\infty(G) \leq RM_1(G) - (\delta - 1)Pl(G).$$

Proof. Let G be any graph. Then,

$$\begin{aligned}
 RM_1(G) &= \sum_{g \sim h} [d(g) + d(h)] \\
 &\leq \sum_{g \sim h} \max\{d(g), d(h)\} + 2(\Delta - 1) \frac{Pl(G)}{2} \\
 (2.7) \quad &\leq RSO_\infty(G) + (\Delta - 1)Pl(G).
 \end{aligned}$$

$$(2.8) \quad \text{Similarly, } RM_1(G) \geq RSO_\infty(G) + (\delta - 1)Pl(G).$$

From equations (2.7) and (2.8), we get

$$RM_1(G) - (\Delta - 1)Pl(G) \leq RSO_\infty(G) \leq RM_1(G) - (\delta - 1)Pl(G)$$

□

Theorem 2.12. Let G be (n, m) -graph with $\Delta \geq 2$. Then

$$\frac{4(\delta - 1)ISI_e(G)}{Pl(G)(\Delta - 1)} \leq RSO_\infty(G) \leq \frac{4(\Delta - 1)ISI_e(G)}{Pl(G)(\delta - 1)}.$$

Equality holds if and only if G is regular.

Proof. Let G be any graph. Then,

$$\begin{aligned}
 ISI_e(G) &= \sum_{g \sim h} \frac{d(g)d(h)}{d(g) + d(h)} \\
 \sum_{g \sim h} \frac{d(g)2(\delta - 1)}{2 \times 2(\Delta - 1)} &\leq \sum_{g \sim h} \frac{d(g)d(h)}{d(g) + d(h)} \leq \sum_{g \sim h} \frac{d(g)2(\Delta - 1)}{2 \times 2(\delta - 1)} \\
 \frac{4(\delta - 1)ISI_e(G)}{Pl(G)(\Delta - 1)} &\leq RSO_\infty(G) \leq \frac{4(\Delta - 1)ISI_e(G)}{Pl(G)(\delta - 1)}.
 \end{aligned}$$

Equality is attained if the graph is regular.

□

3. ENTIRE INF-SOMBOR INDEX

This section contains the precise values of the reformulated Inf-Sombor index for certain graph families.

Observation : Let G be connected graph with $n \geq 3$. Then,

$$(3.1) \quad ESO_\infty(G) = SO_\infty(G) + RSO_\infty(G) + KG_\infty(G).$$

Theorem 3.1. Let G be an r -regular graph with $n \geq 3$ and $r \geq 1$. Then

$$ESO_\infty(G) = \frac{nr^2}{2} + 2n(r - 1)^2 + 2nr(r - 1).$$

Proof. From Theorem 3.3, we have

$$ESO_\infty(G) = SO_\infty(G) + RSO_\infty(G) + 2Pl(G)$$

$$ESO_\infty(G) = \frac{nr^2}{2} + 2n(r - 1)^2 + 2nr(r - 1).$$

□

Theorem 3.2. (1) For complete graph K_n with $n \geq 3$, $ESO_\infty(K_n) = \frac{(n-1)^2n}{2} + n(n - 2)^2(n - 1) + 2n(n - 1)(n - 2)$.

- (2) For cycle C_n with $n \geq 3$, $ESO_\infty(C_n) = 8n$.
 (3) For complete bipartite graph $K_{m,n}$ with $1 < m \leq n$, $ESO_\infty(K_{m,n}) = \max\{m, n\} \cdot mn + \frac{(m+n-2)^2 mn}{2} + 2(m+n-2)mn$.
 (4) For path P_n with $n \geq 3$, $ESO_\infty(P_n) = 4(2n-3)$.
 (5) For star S_n with $n \geq 3$, $ESO_\infty(S_n) = 2(n-1)^2 + \frac{(n-2)^2(n-1)}{2} + (n-2)(n-1)$.

Theorem 3.3. Let G be any connected graph without pendant vertices with $n \geq 2$. Then,

$$ESO_\infty(G) = SO_\infty(G) + RSO_\infty(G) + 2Pl(G).$$

Proof. Let G be a graph without pendant vertices. Then $d(u) \geq 2$ for all $u \in V(G)$. If $e = uv$ is any edge then $d(e) \geq d(u)$ and $d(e) \geq d(v)$.

$$(3.2) \quad \text{Hence, } \sum_{ue} \max\{d(u), d(e)\} = 2Pl(G).$$

From equation (3.1), we have $ESO_\infty(G) = SO_\infty(G) + RSO_\infty(G) + \sum_{ue} \max\{d(u), d(e)\}$. Using equation (3.2) in equation (3.1) we get, $ESO_\infty(G) = SO_\infty(G) + RSO_\infty(G) + 2Pl(G)$. \square

Theorem 3.4. Let G be any connected graph of order $n \geq 3$ with \tilde{p} -pendant vertices. Then $ESO_\infty(G) = SO_\infty(G) + RSO_\infty(G) + 2Pl(G) - \tilde{p}$.

Proof. Let $v_1, v_2, \dots, v_{\tilde{p}}$ be pendant vertices of a graph G and $e_1, e_2, \dots, e_{\tilde{p}}$ be the edges incident to $v_1, v_2, \dots, v_{\tilde{p}}$, respectively. Then

$$\begin{aligned} ESO_\infty(G) &= SO_\infty(G) + RSO_\infty(G) + KG_\infty(G) \\ &= SO_\infty(G) + RSO_\infty(G) + \sum_{v \text{ is incident to } e} \max\{d(v), d(e)\} \\ &= SO_\infty(G) + RSO_\infty(G) + \sum_{i=1}^{\tilde{p}} [\max\{d(v_i), d(e_i)\} + \max\{d(u_i), d(e_i)\}] + \\ &\quad \sum_{e_i \in E(G)/\{e_1, e_2, \dots, e_{\tilde{p}}\}} [\max\{d(v_i), d(e_i)\} + \max\{d(u_i), d(e_i)\}] \\ &= SO_\infty(G) + RSO_\infty(G) + d(e_1) + d(e_1) - 1 + d(e_2) + d(e_2) - 1 + \dots + d(e_{\tilde{p}}) + d(e_{\tilde{p}}) - 1 + \\ &\quad 2d(e_{\tilde{p}+1}) + \dots + 2d(e_m) \\ ESO_\infty(G) &= SO_\infty(G) + RSO_\infty(G) + 2Pl(G) - \tilde{p}. \end{aligned}$$

\square

3.1. Inequalities related to Entire Inf-Sombor index.

Theorem 3.5. For any graph G with $\Delta \geq 2$,

$$\frac{EM_2(G)}{2(\Delta-1)} \leq ESO_\infty(G) \leq \frac{EM_2(G)}{\delta}.$$

Proof. Let G be any graph. Then

$$\begin{aligned}
 ESO_{\infty}(G) &= \sum_{\{gh\} \subseteq B(G)} \max\{d(g), d(h)\} \\
 &= \sum_{\{gh\} \subseteq B(G)} \frac{\max\{d(g), d(h)\}d(x)d(y)}{d(x)d(y)} \\
 &\leq \frac{EM_2(G)}{\delta}
 \end{aligned}$$

Similarly, $ESO_{\infty}(G) = \sum_{\{gh\} \subseteq B(G)} \max\{d(g), d(h)\}$

$$\begin{aligned}
 &= \sum_{\{gh\} \subseteq B(G)} \frac{\max\{d(g), d(h)\}d(x)d(y)}{d(x)d(y)} \\
 &\geq \frac{EM_2(G)}{2(\Delta - 1)}
 \end{aligned}$$

$$\frac{EM_2(G)}{2(\Delta - 1)} \leq ESO_{\infty}(G) \leq \frac{EM_2(G)}{\delta}.$$

□

Theorem 3.6. For any graph G with $\Delta \geq 2$,

$$\frac{\delta}{8(\Delta - 1)^2} EF(G) \leq ESO_{\infty}(G) \leq \frac{\Delta - 1}{\delta^2} EF(G).$$

Proof. Let G be any graph. Then

$$\begin{aligned}
 ESO_{\infty}(G) &= \sum_{\{g,h\} \subseteq B(G)} \max\{d(g), d(h)\} \\
 &= \sum_{\{g,h\} \subseteq B(G)} \frac{\max\{d(g), d(h)\}[d(x)^2 + d(y)^2]}{[d(x)^2 + d(y)^2]} \\
 (3.3) \quad &\leq \frac{(\Delta - 1)}{\delta^2} EF(G)
 \end{aligned}$$

$$\begin{aligned}
 \text{Similarly, } ESO_{\infty}(G) &= \sum_{\{g,h\} \subseteq B(G)} \frac{\max\{d(g), d(h)\}[d(x)^2 + d(y)^2]}{[d(x)^2 + d(y)^2]} \\
 &\geq \sum_{\{g,h\} \subseteq B(G)} \frac{[d(x)^2 + d(y)^2]\delta}{2 \cdot 4(\Delta - 1)^2} \\
 (3.4) \quad &\geq \frac{\delta}{8(\Delta - 1)^2} EF(G)
 \end{aligned}$$

From equations (3.3) and (3.4), we get

$$\frac{\delta}{8(\Delta - 1)^2} EF(G) \leq ESO_{\infty}(G) \leq \frac{\Delta - 1}{\delta^2} EF(G).$$

□

Theorem 3.7. For any graph G , $\delta^2 ER(G) \leq ESO_{\infty}(G) \leq 4(\Delta(G) - 1)^2 ER(G)$.




FIGURE 1. Regular Dendrimers

Proof. Let G be any graph. Then

$$\begin{aligned}
 ESO_{\infty}(G) &= \sum_{\{g,h\} \subseteq B(G)} \max\{d(g), d(h)\} \\
 &= \sum_{\{g,h\} \subseteq B(G)} \frac{\max\{d(g), d(h)\} \sqrt{d(x)d(y)}}{\sqrt{d(x)d(y)}} \\
 &\leq \sum_{\{g,h\} \subseteq B(G)} \frac{\sqrt{4(\Delta-1)^2 \cdot 2(\Delta-1)}}{\sqrt{d(x)d(y)}} \\
 (3.5) \quad &\leq 4(\Delta-1)^2 ER(G).
 \end{aligned}$$

$$(3.6) \quad \text{Similarly, } ESO_{\infty}(G) = \sum_{\{g,h\} \subseteq B(G)} \max\{d(g), d(h)\} \geq \delta^2 ER(G)$$

(3.5) and (3.6), we get $\delta^2 ER(G) \leq ESO_{\infty}(G) \leq 4(\Delta-1)^2 ER(G)$. \square

4. RESULTS ON DENDRIMERS

One of the most significant hyperbranched nanostructures is the dendrimer, which can be synthesized by convergent or divergent processes and assembled from monomers via a nanoscale production process. Nanotubes, nanolatex, chemical sensors, colored glass, micro and macro capsules, and photon funnels, such as artificial antennas are all made of dendrimers. [30, 31, 33, 37, 38, 39, 40, 41]

4.1. Regular dendrimer. Regular dendrimers, which are highly branched and scattered macro molecules, are a unique type of polymeric material found in chemical trees because of the significant influence they have on the physical and chemical properties of the molecules. There is a central vertex (v) in a regular dendrimer tree ($T_{k,d}$). Let the growth of the regular dendrimer be k i.e the distance between the central vertex to the pendant vertex, and each nonpendant vertex of $T_{k,d}$ has a degree of $d \geq 2$. Figure 1 shows examples of dendrimers $T_{3,4}$.

Lemma 4.1. [32] *If $T_{k,d}$ is a tree with central vertex v , then*

- (1) $T_{k,d}$ has d branches, $\frac{(d-1)^k-1}{d-2}$ vertices, $\frac{(d-1)^{k-1}-1}{d-2}$ nonpendant vertices, $(d-1)^{k-1}$ pendant vertices.
- (2) The order of $T_{k,d}$ is $1 + \frac{((d-1)^k-1)d}{d-2}$.




FIGURE 2. The particular values of the indices associated with the Inf-Sombor index

TABLE 2. The calculated values of the indices associated with the Inf-Sombor index

Indices	Calculated values
$Pl(T_{k,d})$	$\left[1 + \frac{d[(d-1)^k-1]}{d-2} - d(d-1)^{k-1}\right] d(d-1)$
$SO_{\infty}(T_{k,d})$	$\frac{(d-1)^k d^2 - d^2}{d-2}$
$RSO_{\infty}(T_{k,d})$	$2(d-1) \left\{ \frac{d(d-1)}{2} \left[\frac{d[(d-1)^k-1]}{d-2} - d(d-1)^{k-1} + 1 \right] - \frac{d(d-1)^{k-1}(d-2)}{2} \right\}$
$ESO_{\infty}(T_{k,d})$	$\frac{d^2(d-1)^k - d^2}{d-2} + 2(d-1) \left\{ \frac{d(d-1)}{2} \left[1 + \frac{d[(d-1)^k-1]}{d-2} - d(d-1)^{k-1} \right] - \frac{d(d-1)^{k-1}(d-2)}{2} \right\}$ $+ 2d(d-1) \left[1 + \frac{d[(d-1)^k-1]}{d-2} - d(d-1)^{k-1} \right] - d(d-1)^{k-1}$

TABLE 3. The particular values of the indices associated with Inf-Sombor index

Growth k	Indices associated with Inf-Sombor related indices		
	$SO_{\infty}(T_{k,d})$	$RSO_{\infty}(T_{k,d})$	$ESO_{\infty}(T_{k,d})$
k=1	16	12	52
k=2	64	108	292
k=3	208	180	796
k=4	1280	1260	3812

Mathematically, the calculated values of the indices associated with the Inf-Sombor index are given in Table 2. We calculated particular values of the indices associated with the Inf-Sombor index of different growth k for $1 \leq k \leq 4$ as shown in Table 3 and its graphical comparison is shown in Figure 2.

4.2. Polyamidoamine (PAMAM), Polypropylenimine (PPI) and Polypropyleneamine (POPAM) Dendrimers. PAMAM dendrimers are characterized by their size, shape, and multifunctional terminal surface. They are hyperbranched dendrimers with a restricted molecular weight distribution and unmatched molecular homogeneity. It seems that the polypropylenimine dendrimers are appealing nonviral vectors for antisense oligonucleotides, small interfering RNA, and gene delivery. On the other hand, little is known about how PPI synthetic gene delivery vectors affect global gene expression. The Polypropyleneamine dendrimers closely resemble PPI dendrimers.

We have calculated the indices related to inf-Sombor index in table 4 and compared them graphically for growth values $k = 1, 2, 3, \text{ and } 4$, which is shown in Figure 4a, 4b, 4c and 4d

TABLE 4. The calculated values of the indices associated with the different dendrimers


Dendrimers	Indices	Calculated values
D_1	$Pl(D_1)$	$114 \times 2^k - 60$
	$SO_\infty(D_1)$	$69 \times 2^k - 33$
	$RSO_\infty(D_1)$	$162 \times 2^k - 87$
	$ESO_\infty(D_1)$	$450 \times 2^k - 237$
D_2	$Pl(D_2)$	$152 \times 2^k - 72$
	$SO_\infty(D_2)$	$164 \times 2^k - 76$
	$RSO_\infty(D_2)$	$216 \times 2^k - 104$
	$ESO_\infty(D_2)$	$672 \times 2^k - 320$
D_3	$Pl(D_3)$	$76 \times 2^k - 72$
	$SO_\infty(D_3)$	$82 \times 2^k - 76$
	$RSO_\infty(D_3)$	$108 \times 2^k - 104$
	$ESO_\infty(D_3)$	$336 \times 2^k - 320$
D_4	$Pl(D_4)$	$36 \times 2^k - 28$
	$SO_\infty(D_4)$	$38 \times 2^k - 28$
	$RSO_\infty(D_4)$	$48 \times 2^k - 40$
	$ESO_\infty(D_4)$	$156 \times 2^k - 124$
D_5	$Pl(D_5)$	$72 \times 2^k - 28$
	$SO_\infty(D_5)$	$52 \times 2^k - 28$
	$RSO_\infty(D_5)$	$168 \times 2^k - 40$
	$ESO_\infty(D_5)$	$360 \times 2^k - 204$

Based on the vertices degrees, we have three partitions of the vertex set: $V_3(D_j) = \{u : d(u) = 3\}$, $V_2(D_j) = \{u : d(u) = 2\}$ and $V_1(D_j) = \{u : d(u) = 1\}$.


Also, the four partitions of the edge set of D_j based on the degrees of end vertices are as follows:

$$\begin{aligned}
E_1(D_j) &= \{e = uv \in E(D_j) : d(v) = 2 \text{ and } d(u) = 1\} \\
E_2(D_j) &= \{e = uv \in E(D_j) : d(v) = 3 \text{ and } d(u) = 1\} \\
E_3(D_j) &= \{e = uv \in E(D_j) : d(v) = 2 \text{ and } d(u) = 2\} \\
E_4(D_j) &= \{e = uv \in E(D_j) : d(v) = 3 \text{ and } d(u) = 2\}.
\end{aligned}$$


Dendrimers	Class Number	Order of vertex and edge class	
D_1	$1 \leq i \leq t$	$ V_i(D_j) $	$ E_i(D_j) $
	$i = 1$	$9 \times 2^k - 3$	3×2^k
	$i = 2$	$30 \times 2^k - 15$	$6 \times 2^k - 3$
	$i = 3$	$9 \times 2^k - 5$	$18 \times 2^k - 9$
	$i = 4$	-	$21 \times 2^k - 12$
D_2	$i = 1$	$12 \times 2^k - 4$	4×2^k
	$i = 2$	$40 \times 2^k - 18$	$8 \times 2^k - 1$
	$i = 3$	$12 \times 2^k - 6$	$24 \times 2^k - 11$
	$i = 4$	-	$28 \times 2^k - 14$
D_3	$i = 1$	$6 \times 2^k - 4$	2^{k+1}
	$i = 2$	$20 \times 2^k - 18$	$4 \times 2^k - 4$
	$i = 3$	$6 \times 2^k - 6$	$12 \times 2^k - 11$
	$i = 4$	-	$14 \times 2^k - 14$
D_4	$i = 1$	2^{k+1}	2^{k+1}
	$i = 2$	$12 \times 2^k - 8$	-
	$i = 3$	$2 \times 2^k - 2$	$8 \times 2^k - 5$
	$i = 4$	-	$6 \times 2^k - 6$
D_5	$i = 1$	4×2^k	4×2^k
	$i = 2$	$24 \times 2^k - 8$	-
	$i = 3$	$4 \times 2^k - 2$	$16 \times 2^k - 5$
	$i = 4$	-	$12 \times 2^k - 6$

TABLE 5. Order of vertex and edge class of D_j 


(A) Comparison of Platt Number of Dendrimers




(B) Comparison of Infinite Sombor index of Dendrimers




(C) Comparison of Reformulated Inf Sombor index of Dendrimers




(D) Comparison of Entire Inf Sombor index of Dendrimers




(A) First type of polyamidoamine dendrimer D_1




(B) Second type of polyamidoamine dendrimer D_2



(C) First type of polypropylenimine D_3



(D) Polypropylenimine octaamine dendrimer D_4



(E) Polypropyleneamine dendrimer (POPAM) D_5

5. CONCLUSIONS

To achieve more accurate estimates of intermolecular forces, one needs to consider the relations between edges and vertices in addition to the relation between vertices. This is because intermolecular forces occur not just between atoms but also between atoms and bonds. Hence, we studied the edge version of the Inf-Sombor and Entire Sombor indexes and gave some bounds. In the end, we calculated these indices for the different dendrimers and made a graphical comparison. The following can be taken into consideration for further study.

- (1) Characterize these indices with respect to other degree-based topological indices.

(2) Finding extremal graphs for these indices will be an interesting task.

6. DECLARATION

Funding: This research received no external funding.

Conflicts of Interest: The authors declares that there are no conflicts of interest.

Acknowledgement: Grateful thanks to the anonymous referees for their insightful comments and suggestions.

REFERENCES

- [1] I. Gutman, *Geometric approach to degree-based topological indices: Sombor indices*, MATCH-Communications in Mathematical and in Computer Chemistry, **86** (2021), 11-16.
- [2] T. Réti, T. Došlić, and A. Ali, *On the Sombor index of graphs*, Contributions to Mathematics, **3** (2021), 11-18.
- [3] J. L. Gross, J. Yellen and M. Anderson, *Graph Theory and its Applications* Chapman and Hall/CRC, 2018.
- [4] P. Zhang and G. Chartrand, *Introduction to Graph Theory*, **2** Tata McGraw-Hill, New York, 2006.
- [5] I. Gutman, and K. C. Das, *The first Zagreb index 30 years after*, MATCH-Communications in Mathematical and in Computer Chemistry, **50** (2004), 83-92.
- [6] I. Gutman, and N. Trinajstić, *Graph theory and molecular orbitals. Total phi-electron energy of alternate hydrocarbons*, Chemical Physics Letters, **17** (1972), 535-538.
- [7] B. Zhou, *Zagreb indices*, MATCH-Communications in Mathematical and in Computer Chemistry **52** (2004), 113-118.
- [8] B. Furtula and I. Gutman, *A forgotten topological index*, Journal of Mathematical Chemistry, **53** (2015), 1184-1190.
- [9] C. Zhongyuan and C. Zhibo, *Lower and upper bounds of the forgotten topological index*, MATCH Communications in Mathematical and in Computer Chemistry, **76** (2016), 635-648.
- [10] L. Xueliang and Y. Shi, *A survey on the Randic index*, MATCH Communications in Mathematical and in Computer Chemistry, **59** (2008), 127-156.
- [11] M. Randic, *Characterization of molecular branching*, Journal of the American Chemical Society, **97** (1975), 6609-6615.
- [12] J. A. Méndez-Bermúdez, R. Aguilar-Sanchez, E. D. Molina, J. M. Rodríguez *Mean sombor index*, arXiv preprint, (2021).
- [13] J. R. Platt *Prediction of isomeric differences in paraffin properties*, The Journal of Physical Chemistry, **56** (1952), 328-336.
- [14] M. M. Belavadi and T. A. Mangam, *Platt number of total graphs*, International Journal of Applied Mathematics, **31** (2018), 593.
- [15] A. Akbar and D. Dimitrov, *On the extremal graphs with respect to bond incident degree indices*, Discrete Applied Mathematics, **238** (2018), 32-40.
- [16] I. Gutman and E. Estrada, *Topological indices based on the line graph of the molecular graph*, Journal of Chemical Information and Computer Sciences, **36** (1996), 541-543.
- [17] A. Saleh, and I. N. Cangul, *On the entire Randic index of graphs*, Advances in Applied Mathematics, **20** (2021), 1559-1569.
- [18] F. Movahedi and M. H. Akhbari *Entire Sombor index of graphs*, Iranian Journal of Mathematical Chemistry, **14** (2023), 33-45.
- [19] N. Harish and B. Chaluvaram, *Some Inequalities for the Entire Sombor Index*, International Journal of Mathematical Combinatorics, **1** (2024), 50-64.
- [20] N. Harish, B. Sarveshkumar and B. Chaluvaram, *The reformulated Sombor index of a graph*, Transactions on Combinatorics, **13** (2024), 1-16.

- [21] A. Miličević, S. Nikolić and N. Trinajstić, *On reformulated Zagreb indices*, Molecular Diversity, **8** (2004), 393-399.
- [22] V. R. Kulli, N. Harish, B. Chaluvvaraju, I. Gutman, *Mathematical properties of KG Sombor index*, Bulletin of International Mathematical Virtual Institute, **12** (2022), 379-386.
- [23] A. Alwardi, A. Alqesmah, R. Rangarajan and I. N. Cangul, *Entire Zagreb indices of graphs*, Discrete Mathematics, Algorithms and Applications, **10** (2018), 1850037.
- [24] I. Gutman, V. R. Kulli, B. Chaluvvaraju and H. S. Boregowda, *On Banhatti and Zagreb indices*, Journal of International Mathematical Virtual Institute, **7** (2017), 53-67.
- [25] V. R. Kulli, B. Chaluvvaraju and H. S. Boregowda, *The hyper-Zagreb index of some derived graphs*, International Journal of Mathematical Archive, **9** (2018), 124-129.
- [26] H. S. Boregowda, *Topological indices of some derived graphs through M-Polynomials*, Advances and Applications in Mathematical Sciences, (2022).
- [27] H. S. Boregowda, *Wiener Type Indices Of Certain Classes of Trees*, South East Asian Journal of Mathematics and Mathematical Sciences, **17** (2021), 241-250.
- [28] V. R. Kulli, B. Chaluvvaraju and H. S. Boregowda, *Some degree based connectivity indices of Kulli cycle windmill graphs*, South Asian Journal of Mathematics, **6** (2016), 263-268.
- [29] H. Marzieh, *Study of inverse sum indeg index*, Journal of Nanoscience and Nanotechnology, **7** (2017), 103-109.
- [30] B. Klajnert and B. Maria, *Dendrimers: properties and applications*, Acta Biochimica Polonica, **48** (2001), 199-208.
- [31] A. R. Ashrafi and M. Mirzargar, *PI, Szeged and edge Szeged indices of an infinite family of nanostar dendrimers*, Indian Journal of Chemistry, **47** (2008), 538-541.
- [32] Ü. Şener and B. Şahin *Total domination number of regular dendrimer graph*, Turkish Journal of Mathematics and Computer Science, **11** (2019), 81-84.
- [33] V. Mathad, P. Padmapriya and M. A. Sangamesha, *Some topological indices of phosphorus containing dendrimers*, Palestine Journal of Mathematics, **10** (2021), 151-160.
- [34] V. R. Kulli, B. Chaluvvaraju and H. S. Boregowda, *Some bounds of sum connectivity Banhatti index of graphs*, Palestine Journal of Mathematics, **8** (2019), 355-364.
- [35] I. Sarkar, N. Manjunath, B. Chaluvvaraju and V. Loksha *Bounds of Sombor Index for F-Sum Operation*, Palestine Journal of Mathematics, **12** (2023), 504-516.
- [36] A. Jahanbani, *The Multiplicative Hyper-Zagreb index of Graph Operations*, Palestine Journal of Mathematics, **9** (2020), 82-96.
- [37] V. Loksha, S. Jain, M. Muddalapuram, A. S. Çevik and I. N. Cangül, *Bounds for the sum of cubes of vertex degrees of splice graphs*, TWMS Journal Of Applied And Engineering Mathematics, **10** (2020), 741-750.
- [38] P. S. Ranjini, V. Loksha and S. Kumar, *The K. Banhatti Indices of Certain Graphs*, Southeast Asian Bulletin of Mathematics, **46** (2022), 453.
- [39] v. Loksha, P. S. Ranjini and M. R. Harini, *Limits for the Some Topological Indices of Vertex Corona Graph*, International Journal of Mathematical Combinatorics, **4** (2024), 42.
- [40] A. S. Maragadam, K. Nirupadi, D. Dafik, V. Loksha, M. Manjunath and I. H. Agustin, *Neighborhood vertex corona product of graphs in a semi-total point graph*, AIP Conference Proceedings, **3176** (2024).
- [41] S. Jain, A. S. Maragadam, V. Loksha, *Topological Aspects of Synthetic Polymers through NM Polynomials*, 2nd International Conference on Neural Networks and Machine Learning, (2023).