

A COLLOCATION METHOD TO THE SOLUTION OF NONLINEAR FREDHOLM-HAMMERSTEIN INTEGRAL AND INTEGRO-DIFFERENTIAL EQUATIONS

FARSHID MIRZAEI * AND ELHAM HADADIYAN

ABSTRACT. This paper presents a computational technique for the solution of the nonlinear Fredholm-Hammerstein integral and integro-differential equations. A hybrid of block-pulse functions and the second kind Chebyshev polynomials (hereafter called as **HBC**) is used to approximate the nonlinear Fredholm-Hammerstein integral and integro-differential equations. The main properties of HBC are presented. Also, the operational matrix of integration together with the Newton-Cotes nodes are applied to reduce the computation of the nonlinear Fredholm-Hammerstein integral and integro-differential equations into some algebraic equations. The efficiency and accuracy of the proposed method have been shown by three numerical examples.

Key Words: Nonlinear Fredholm-Hammerstein integral equation; Block-pulse function; Second kind Chebyshev polynomial; Collocation method.

2010 Mathematics Subject Classification: 45B05, 45G10, 45J05, 65D30.

1. INTRODUCTION

Integral equations play an important role in many branches of linear and nonlinear functional analysis and their applications in elasticity, engineering, mathematical physics and contact mixed problems [2, 13, 17, 23]. Several methods for solving such equations are available. Typical examples include spectral [18, 21] and transform methods

Received: 30 December 2012, Accepted: 5 March 2013. Communicated by D. Khojasteh Salkuyeh

*Address correspondence to Farshid Mirzaee; E-mail: f.mirzaee@malayeru.ac.ir

© 2013 University of Mohaghegh Ardabili.

[3, 25] as well as other methods [9, 14, 19, 26]. Many methods such as variational iteration [27], homotopy perturbation [8, 22], domain decomposition [5, 12], RBF [10, 11], wavelets [16, 29, 31] and others [1] have been developed and improved to obtain numerical solutions.

In the present paper, we consider application of HBC to obtain the numerical solution of the nonlinear Fredholm-Hammerstein integral equation:

$$(1.1) \quad y(t) = x(t) + \lambda \int_0^1 k(t, s)F(y(s))ds, \quad 0 \leq t < 1 \quad ,$$

and the nonlinear Fredholm-Hammerstein integro-differential equation:

$$(1.2) \quad \begin{cases} y'(t) = x(t) + \lambda \int_0^1 k(t, s)F(y(s))ds \\ y(0) = y_0, \quad 0 \leq t < 1, \end{cases}$$

where λ and y_0 are constants and $x(t)$ and $k(t, s)$ are assumed to be in $L_2[0, 1)$ and $L_2([0, 1) \times [0, 1))$, respectively. We assume that Eqs. (1.1) and (1.2) have a unique solution $y(t)$ as a linear combination of the HBC with some unknown coefficients [15]. The properties of HBC together with the Newton-Cotes nodes [24] are then applied to evaluate the unknown coefficients and find an approximate solution to Eqs. (1.1) and (1.2).

This paper is organized as follows: In Section 2, we describe the basic formulation of HBC required for our subsequent development. Sections 3 and 4 are devoted to the solution of Eqs. (1.1-1.2) by using HBC. In Section 5, we report our numerical results and demonstrate the accuracy of the proposed scheme by considering the numerical examples.

2. PROPERTIES OF HBC

2.1. HBC Definition. A set of block-pulse functions $\phi_n(t), n = 1, 2, \dots, N$, on interval $[0, 1)$ is defined as (see [21]):

$$(2.1) \quad \phi_n(t) = \begin{cases} 1, & \frac{n-1}{N} \leq t < \frac{n}{N} \\ 0, & \text{otherwise} \end{cases} ,$$

where N is an arbitrary positive integer.

The m th second kind Chebyshev polynomial in interval $[-1, 1]$ is defined as follows (see [25]):

$$(2.2) \quad S_m(t) = \frac{\sin((m+1) \arccos(t))}{\sqrt{1-t^2}} .$$

Also, we can, obtain the second kind Chebyshev polynomials by the following recursive formulas:

$$(2.3) \quad \begin{cases} S_0(t) = 1, & S_1(t) = 2t, \\ S_{m+1}(t) = 2tS_m(t) - S_{m-1}(t), & m = 1, 2, \dots \end{cases} .$$

Further, HBC $H_{nm}(t), n = 1, 2, \dots, N, m = 0, 1, \dots, M - 1$, on the interval $[0, 1)$ are defined as:

$$(2.4) \quad H_{nm}(t) = \phi_n(t)S_m(2Nt - 2n + 1).$$

2.2. Function approximation. A function $f(t) \in L_2[0, 1)$ may be expanded as

$$(2.5) \quad f(t) \approx \sum_{n=1}^N \sum_{m=0}^{M-1} f_{nm}H_{nm}(t),$$

where

$$(2.6) \quad f_{nm} = \frac{2}{\pi} \int_{-1}^1 f\left(\frac{t+2n-1}{2N}\right)S_m(t)\sqrt{1-t^2}dt.$$

We can rewrite $f(t)$ as:

$$(2.7) \quad f(t) \approx \sum_{n=1}^N F_n H_k(t) = FH(t),$$

where

$$(2.8) \quad F_n = [f_{n0}, \dots, f_{n(M-1)}], \quad F = [F_1, \dots, F_N],$$

and

$$(2.9) \quad H_n(t) = [H_{n0}(t), \dots, H_{n(M-1)}(t)]^T, \quad H(t) = [H_1^T(t), \dots, H_N^T(t)]^T.$$

2.3. The operational matrix of integration. The integral part in Eq. (2.9) is given by:

$$(2.10) \quad \int_0^t H(\tau)d\tau \approx PH(t),$$

where

$$(2.11) \quad P = \frac{1}{2N}I \otimes \hat{P} + \sum_{n=1}^{N-1} \sum_{i=1}^{N-n} \frac{1}{N} E_{i(i+n)}^{(N)} \otimes \sum_{n=1}^{\lfloor \frac{M+1}{2} \rfloor} \frac{1}{2n-1} E_{(2n-1)1}^{(M)},$$

$$(2.12) \quad \hat{P} = E_{11}^{(M)} - \frac{3}{4}E_{21}^{(M)} + \frac{1}{2} \left(\sum_{n=1}^{M-1} \frac{1}{n} E_{n(n+1)}^{(M)} - \sum_{n=2}^{M-1} \frac{1}{n+1} E_{(n+1)n}^{(M)} \right) + \sum_{n=3}^M \frac{(-1)^{n-1}}{n} E_{n1}^{(M)}.$$

Where $E_{ij}^{(m)}$ is the $m \times m$ matrix with entry 1 at (i, j) and zeros elsewhere and $\lceil \frac{M+1}{2} \rceil$ is the greatest integer part of $\frac{M+1}{2}$.

For more details, see [30].

In Eq. (2.11), notation \otimes denotes Kronecker product which defined as:

$$A_{n \times p} \otimes B_{m \times q} = \begin{bmatrix} a_{11}B & a_{12}B & \dots & a_{1p}B \\ a_{21}B & a_{22}B & \dots & a_{2p}B \\ \vdots & \vdots & \vdots & \vdots \\ a_{n1}B & a_{n2}B & \dots & a_{np}B \end{bmatrix}_{nm \times pq}.$$

2.4. Expression of producing functions. Let $g(t), f(t) \in L_2[0, 1)$, then the expressions of $g(t)$ and $f(t)$ are defined as:

$$(2.13) \quad g(t) \approx \sum_{n=1}^N \sum_{m=0}^{M-1} g_{nm} H_{nm}(t), \quad f(t) \approx \sum_{n=1}^N \sum_{m=0}^{M-1} f_{nm} H_{nm}(t).$$

Then

$$(2.14) \quad g(t)f(t) \approx \sum_{n=1}^N \sum_{i=0}^{M-1} \sum_{j=0}^{M-1} g_{ni} f_{nj} H_{ni}(t) H_{nj}(t).$$

From

$$(2.15) \quad H_{ni}(t) H_{nj}(t) \approx \sum_{m=0}^{M-1} d_{nm}^{(i,j)} H_{nm}(t),$$

where

$$(2.16) \quad d_{nm}^{(i,j)} = \frac{2}{\pi} \int_{-1}^1 S_i(t) S_j(t) S_m(t) \sqrt{1-t^2} dt,$$

we have:

$$(2.17) \quad g(t)f(t) \approx \sum_{n=1}^N \sum_{m=0}^{M-1} \left(\sum_{i=0}^{M-1} \sum_{j=0}^{M-1} d_{nm}^{(i,j)} g_{ni} f_{nj} \right) H_{nm}(t) = \sum_{n=1}^N \sum_{m=0}^{M-1} \tilde{g}_{nm} H_{nm}(t),$$

whereas

(2.18)

$$\tilde{g}_{nm} = \sum_{i=0}^{M-1} \sum_{j=0}^{M-1} d_{nm}^{(i,j)} g_{ni} f_{nj} = \hat{g}_{nm} F_n^T, \quad F_n = [f_{n0}, \dots, f_{n(M-1)}],$$

and

$$(2.19) \quad \hat{g}_{nm} = \left[\sum_{i=0}^{M-1} d_{nm}^{(i,0)} g_{ni}, \dots, \sum_{i=0}^{M-1} d_{nm}^{(i,M-1)} g_{ni} \right].$$

Therefore, using Eq. (2.5), it may be assumed that:

$$(2.20) \quad k(t, s)w(s) \approx \sum_{n=1}^N \sum_{m=0}^{M-1} \tilde{k}_{nm}(t) H_{nm}(s),$$

where

$$(2.21) \quad w(s) \approx WH(s),$$

$W = [W_1, \dots, W_N]$ and $H(s)$ are given in Eq. (2.8) and Eq. (2.9), respectively, and

$$(2.22) \quad \tilde{k}_{nm}(t) = \hat{k}_{nm}(t) W_n^T,$$

$$(2.23) \quad \hat{k}_{nm}(t) = \left[\sum_{i=0}^{M-1} d_{nm}^{(i,0)} k_{ni}(t), \dots, \sum_{i=0}^{M-1} d_{nm}^{(i,M-1)} k_{ni}(t) \right],$$

$$(2.24) \quad k_{ni}(t) = \frac{2}{\pi} \int_{-1}^1 k(t, \frac{s+2n-1}{2N}) S_i(s) \sqrt{1-s^2} ds.$$

Let

$$(2.25) \quad v(t) = \int_0^1 k(t, s)w(s) ds \approx \sum_{n=1}^N \sum_{m=0}^{M-1} v_{nm} H_{nm}(t),$$

$$(2.26) \quad \hat{k}_{nm}(t) \approx \sum_{j=1}^N \sum_{l=0}^{M-1} \hat{k}_{nm}^{(j,l)} H_{jl}(t),$$

where

$$(2.27) \quad \hat{k}_{nm}^{(j,l)} = \left[\sum_{i=0}^{M-1} d_{nm}^{(i,0)} k_{ni}^{(j,l)}, \dots, \sum_{i=0}^{M-1} d_{nm}^{(i,M-1)} k_{ni}^{(j,l)} \right],$$

$$(2.28) \quad k_{ni}^{(j,l)} = \frac{2}{\pi} \int_{-1}^1 k_{ni} \left(\frac{t+2j-1}{2N} \right) S_l(t) \sqrt{1-t^2} dt.$$

By Eqs. (2.20) and (2.22) we have:

$$(2.29) \quad v(t) \approx \sum_{j=1}^N \sum_{l=0}^{M-1} \sum_{n=1}^N \sum_{m=0}^{M-1} \widehat{k}_{nm}^{(j,l)} \left(\int_{\frac{n-1}{N}}^{\frac{n}{N}} H_{nm}(t) dt \right) W_n^T H_{jl}(t).$$

Hence

$$(2.30) \quad V = \sum_{n=1}^{\lfloor \frac{M+1}{2} \rfloor} \sum_{i=0}^N \sum_{j=1}^N E_{ij}^{(N)} \otimes \frac{1}{N(2n-1)} \widehat{k}_{j(2n-2)}^{(i)} W^T,$$

where

$$(2.31) \quad \widehat{k}_{j(2n-2)}^{(i)} = \left[\widehat{k}_{j(2n-2)}^{T(i,0)}, \dots, \widehat{k}_{j(2n-2)}^{T(i,M-1)} \right]^T.$$

Therefore

$$(2.32) \quad v(t) = \int_0^1 k(t,s)w(s)ds \approx W \widehat{V}^T H(t),$$

where

$$(2.33) \quad \widehat{V} = \sum_{n=1}^{\lfloor \frac{M+1}{2} \rfloor} \sum_{i=0}^N \sum_{j=1}^N E_{ij}^{(N)} \otimes \frac{1}{N(2n-1)} \widehat{k}_{j(2n-2)}^{(i)}.$$

3. NONLINEAR FREDHOLM-HAMMERSTEIN INTEGRAL EQUATION

In this section we consider the Fredholm-Hammerstein integral Eq. (1.1). To solve this equation, we first consider the approximation:

$$(3.1) \quad w(t) = F(y(t)), \quad 0 \leq t < 1,$$

and then substitute it in Eq. (1.1).

From Eq. (1.1) we get

$$(3.2) \quad w(t) = F \left(x(t) + \lambda \int_0^1 k(t,s)w(s)ds \right).$$

By some manipulations as before, we have:

$$(3.3) \quad WH(t) = F(x(t) + \lambda W \widehat{V}^T H(t)).$$

Now, using Newton-Cotes nodes as

$$t_i = \frac{2i-1}{2NM}, \quad i = 1, 2, \dots, NM,$$

we have

$$(3.4) \quad WH(t_i) = F(x(t_i) + \lambda W \widehat{V}^T H(t_i)), \quad i = 1, 2, \dots, NM.$$

By solving this system of nonlinear equations, we find unknown vector W which has NM components. Substituting Eq. (3.1) into Eq. (1.1), yields:

$$(3.5) \quad y(t) = x(t) + \lambda \int_0^1 k(t, s)w(s)ds,$$

and applying Eq. (2.32) to Eq. (3.5), we get

$$(3.6) \quad y(t) = x(t) + \lambda W \widehat{V}^T H(t).$$

In other words, we could find the unknown function $y(t)$.

4. NONLINEAR FREDHOLM-HAMMERSTEIN INTEGRO-DIFFERENTIAL EQUATION

Here, the Fredholm-Hammerstein integro-differential Eq. (1.2) is considered.

In order to solve this equation, we suppose that:

$$(4.1) \quad w(t) = F(y(t)), \quad 0 \leq t < 1,$$

when Eq. (4.1) implies that:

$$(4.2) \quad y(t) = y(0) + \int_0^t y'(\tau)d\tau.$$

We approximate $x(\tau)$ by Eq. (2.7) as follows:

$$(4.3) \quad x(\tau) = XH(\tau),$$

where X and $H(\tau)$ are given in Eq. (2.8) and Eq. (2.9), respectively. Consequently, we have:

$$(4.4) \quad y(t) = y(0) + \int_0^t x(\tau)d\tau + \lambda \int_0^t \int_0^1 k(\tau, s)w(s)dsd\tau \\ \approx y(0) + XPH(t) + \lambda W \widehat{V}^T PH(t),$$

and

$$(4.5) \quad WH(t) = F(y(0) + XPH(t) + \lambda W \widehat{V}^T PH(t)).$$

In order to construct the appropriate approximations for $w(t)$, we collocate Eq. (4.5) in NM points. Now, using Newton-Cotes nodes, we have:

$$(4.6) \quad WH(t_i) = F(y(0) + XPH(t_i) + \lambda W\widehat{V}^T PH(t_i)), \quad i = 1, 2, \dots, NM.$$

Finally, W is obtained by solving Eq. (4.6).

5. NUMERICAL EXAMPLES

To demonstrate the efficiency of our scheme, some numerical examples are presented in this section. The computations associated with the examples were performed by Matlab software on a personal computer.

Example 1. Consider the following nonlinear Fredholm integral equation [20]:

$$y(t) = \cos(t) - 0.4958t + \int_0^1 ts \tan(y(s)) ds, \quad 0 \leq t \leq 1.$$

Taking $N = 1, M = 2$, we obtain the following approximate solution:

$$y_{N,M}(t) = y_{1,2}(t) \approx \cos(t) - 0.4958t + .278886658H_{10}(t) + 0.13944333H_{11}(t),$$

and for $N = 2, M = 4$, we obtain the approximate solution as:

$$y_{N,M}(t) = y_{2,4}(t) \approx \cos(t) - 0.4958t + 0.12394656H_{10}(t) + 0.06197328H_{11}(t) \\ + 0.037183967H_{20}(t) + 0.06197328H_{21}(t).$$

Taking $N = 1, M = 2$ and $N = 2, M = 4$ the solutions are compared with the exact solution $y(t) = \cos(t)$ as presented in Table 1 and depicted graphically in Figure 1.


The 2-norm errors for the proposed method are in agreement with those of the methods of Babolian *et al.* [6] and of Maleknejad *et al.* [20] as seen in Table 2.

Table 1: Numerical results of Example 1 with HBC

Nodes t	Approximate HBC for $N = 1$ and $M = 2$	Error HBC for $N = 1$ and $M = 2$	Approximate HBC for $N = 2$ and $M = 4$	Error HBC for $N = 2$ and $M = 4$
t = 0.0	1.00000000e-0	0.00000000e-0	1.00000000e-0	0.00000000e-0
t = 0.1	1.00120150e-0	6.19733165e-3	9.95002788e-1	1.37700726e-6
t = 0.2	9.92461241e-1	1.23946633e-2	9.80063824e-1	2.75401453e-6
t = 0.3	9.73928484e-1	1.85919950e-2	9.55332358e-1	4.13102179e-6
t = 0.4	9.45850321e-1	2.47893266e-2	9.21055486e-1	5.50802905e-6
t = 0.5	9.08569220e-1	3.09866583e-2	8.77575677e-1	6.88503632e-6
t = 0.6	8.62519605e-1	3.71839899e-2	8.25327353e-1	8.26204358e-6
t = 0.7	8.08223509e-1	4.33813216e-2	7.64832548e-1	9.63905084e-6
t = 0.8	7.46285363e-1	4.95786532e-3	6.96695693e-1	1.10160581e-5
t = 0.9	6.77385953e-1	5.57759849e-3	6.21597575e-1	1.23930654e-5

Table 2: Approximate norm-2 of absolute error for Example 1

Methods	$\ y(t) - y^*(t)\ $
Method of [6]	
m = 2	2.27187e-1
m = 3	1.34023e-1
m = 4	1.07511e-1
Method of [20]	
$j_u = 3$	3.54072e-5
$j_u = 4$	5.61248e-6
$j_u = 5$	7.83725e-7
Present method	
N = 1 , M = 2	7.11353e-2
N = 2 , M = 4	2.31441e-5
N = 3 , M = 4	6.84629e-8



Example 2. Consider the following nonlinear Fredholm integro-differential equation [4]:

$$y'(t) = 1 - \frac{1}{4}t + \int_0^1 tsy^2(s)ds, \quad y(0) = 0.$$

Taking $N = 1, M = 2$, we obtain the following approximate solution:

$$y_{N,M}(t) = y_{1,2}(t) \approx 0.494193699H_{10}(t) + 0.245354959H_{11}(t),$$

and for $N = 2, M = 4$, we obtain the approximate solution as:

$$y_{N,M}(t) = y_{2,4}(t) \approx 0.25H_{10}(t) + 0.125H_{11}(t) - 0.888595270e^{-31}H_{12}(t) \\ + 0.75H_{20}(t) + 0.125H_{21}(t) - 0.888595270e^{-31}H_{22}(t).$$

The obtained solutions for $N = 1, M = 2$ and for $N = 2, M = 4$ are comparable with the exact solution $y(t) = t$ as have shown in Table 3 and Figure 2.

The 2–norm errors of the proposed method are in a good agreement with those of the methods of Babolian *et al.* [6] and of Saeedi *et al.* [28] as seen in Table 4.

Table 3: Numerical results of Example 2 with HBC

Nodes t	Approximate HBC for $N = 1$ and $M = 2$	Error HBC for $N = 1$ and $M = 2$	Error HBC for $N = 2$ and $M = 4$
t = 0.0	3.48378083e-3	3.48378083e-3	0.00000000e-00
t = 0.1	1.01625764e-1	1.62576438e-3	5.68700973e-32
t = 0.2	1.99767748e-1	2.32252056e-4	2.27480389e-31
t = 0.3	2.97909731e-1	2.09026850e-3	5.11830876e-31
t = 0.4	3.96051715e-1	3.94828495e-3	9.09921557e-31
t = 0.5	4.94193699e-1	5.80630139e-3	1.42175243e-30
t = 0.6	5.92335682e-1	7.66431784e-3	2.04732350e-30
t = 0.7	6.90477666e-1	9.52233428e-3	2.78663477e-30
t = 0.8	7.88619649e-1	1.13803507e-2	3.63968623e-30
t = 0.9	8.86761633e-1	1.32383672e-2	4.60647788e-30




Table 4: Approximate norm-2 of absolute error for Example 2

Methods	$\ y(t) - y^*(t)\ $
Method of [6]	
m = 2	4.7434e-01
m = 3	3.0732e-01
m = 4	2.3717e-01
Method of [28]	
K = 2 , M = 1	2.7133e-03
K = 3 , M = 1	6.8179e-04
K = 4 , M = 1	1.6745e-05
Present method	
N = 1 , M = 2	2.2862e-02
N = 2 , M = 4	7.0420e-30
N = 3 , M = 4	3.4729e-38

Example 3. Consider the nonlinear Fredholm integro-differential equation [7]:

$$y'(t) = 2t + \frac{1}{8}(-\pi + \log(4)) + \int_0^1 s \arctan(y(s)) ds, \quad y(0) = 0.$$

Taking $N = 1, M = 2$, we obtain the following approximate solution:

$$y_{N,M}(t) = y_{1,2}(t) \approx 0.299583352H_{10}(t) + 0.243516761H_{11}(t),$$

and for $N = 2, M = 4$, we obtain the following approximate solution:

$$y_{N,M}(t) = y_{2,4}(t) \approx 0.078131491H_{10}(t) + 0.062503245H_{11}(t) + 0.015625H_{12}(t) \\ + 0.578144472H_{20}(t) + 0.187503245H_{21}(t) - 0.015625H_{22}(t).$$


The exact solution of this example is $y(t) = t^2$. As we can see from Table 5 errors for $N = 1, M = 2$ and $N = 2, M = 4$ are acceptable. Also, Figure 3 has a good agreement with the exact solution. The 2-norm errors of the proposed method are in a good agreement with those of the methods of Babolian *et al.* [6] and of Berenguer *et al.* [7] as seen in Table 6.

Table 5: Numerical results of Example 3 with HBC

Nodes t	Approximate HBC for $N = 1$ and $M = 2$	Error HBC for $N = 1$ and $M = 2$	Approximate HBC for $N = 2$ and $M = 4$	Error HBC for $N = 2$ and $M = 4$
t = 0.0	1.87500000e-1	1.87500000e-1	0.00000000e-0	0.00000000e-0
t = 0.1	9.00833296e-2	1.00083330e-1	1.00025963e-2	2.59632768e-6
t = 0.2	7.33334086e-2	3.26666591e-2	4.00051927e-2	5.19265536e-6
t = 0.3	1.04750011e-1	1.47500113e-2	9.00077890e-2	7.78898304e-6
t = 0.4	2.02121667e-1	4.21666817e-2	1.60010385e-1	1.03853107e-5
t = 0.5	2.99583352e-1	4.95833521e-2	2.50012982e-1	1.29816384e-5
t = 0.6	3.97000023e-1	3.70000226e-2	3.60015578e-1	1.55779661e-5
t = 0.7	4.94416693e-1	4.41669301e-3	4.90018174e-1	1.81742938e-5
t = 0.8	5.91833363e-1	4.81666366e-2	6.40020771e-1	2.07706214e-5
t = 0.9	6.89250034e-1	1.20749966e-1	8.10023367e-1	2.33669491e-5

Table 6: Approximate norm-2 of absolute error for Example 3

Methods	$\ y(t) - y^*(t)\ $
Method of [6]	
m = 2	3.40827-1
m = 3	2.43755e-1
Method of [7]	
j = 9 , h = 6	5.42945e-3
j = 17 , h = 6	1.53327e-3
Present method	
N = 1 , M = 2	2.62647e-1
N = 2 , M = 4	4.38311e-5



6. CONCLUSION

In this paper, HBC method has successfully applied to compute the approximate solution of certain nonlinear Fredholm-Hammerstein integral and integro-differential equations. The accuracy and applicability of the method were investigated through some numerical examples. The numerical results showed that the accuracy of the obtained solutions was satisfactory. Furthermore, the current method can be used by increasing N and M until the results reach an appropriate accuracy.

This method can be easily extended and applied to a system of nonlinear Fredholm-Hammerstein integral and integro-differential equations.

REFERENCES

- [1] S. Abbasbandy, *Numerical solution of the integral equations: Homotopy perturbation method and Adomian's decomposition method*, Appl. Math. Comput., **173** (2006) 493–500.
- [2] M.A. Abdou, A.A. Badr, *On a method for solving an integral equation in the displacement contact problem*, J. Appl. Math. Comput., **127** (2002) 65–78.
- [3] A. Arikoglu, I. Ozkol, *Solutions of integral and integro-differential equations systems by using differential transform method*, Comput. Math. Appl., **56** (2008) 2411–2417.
- [4] Z. Avazzadeh, M. Heydari, G.B. Loghmani, *Numerical solution of Fredholm integral equations of the second kind by using integral mean value theorem*, Appl. Math. Model., **35** (2011) 2374–2383.
- [5] E. Babolian, J. Biazar, A.R. Vahidi, *The decomposition method applied to systems for Fredholm integral equations of the second kind*, Appl. Math. Comput., **148** (2004) 443–452.
- [6] E. Babolian, K. Maleknejad, M. Mordad, B. Rahimi, *A numerical method for solving Fredholm-Volterra integral equations in two-dimensional spaces using block-pulse functions and an operational matrix*, J. Comput. Appl. Math., **235** (2011) 3965–3971.
- [7] M. I. Berenguer, M. V. Fernandez Munoz, A. I. Garralda-Guillem, M. Ruiz Galan, *A sequential approach for solving the Fredholm integro-differential equation*, Appl. Numer. Math., (2011) In Press.
- [8] J. Biazar, H. ghazvini, *Numerical solution for special nonlinear Fredholm integral equation by HPM*, Appl. Math. Comput., **195** (2008) 681–687.
- [9] P. Bräuer, M. Fassler, M. Jaroniec, *Numerical solutions of the adsorption integral equation utilizing the spline functions*, Thin Solid Films, **123** (1985) 245–272.
- [10] A. Golbabai, M. Mammadov, S. Seifollahi, *Solving a system of nonlinear integral equations by an RBF network*, Comput. Math. Appl., **57** (2009) 1651–1658.
- [11] A. Golbabai, S. Seifollahi, *Numerical solution of the second kind integral equations using radial basis function networks*, Appl. Math. Comput., **174** (2006) 877–883.

- [12] A. Gorguis, *Charpit and Adomian for solving integral equations*, Appl. Math. Comput., **193** (2007) 446–454.
- [13] R. Grimmer, J.H. Liu, *Singular perturbations in viscoelasticity*, Rocky Mountain J. Math., **24** (1994) 61–75.
- [14] M. Inoue, M. Kuroumaru, S. Yamaguchi, *A solution of Fredholm integral equation by means of the soline fit approximation*, Comput. Fluids, **7** (1979) 33–46.
- [15] Z. H. Jiang, W. Schaufelberger, *Block Pulse functions and their applications in control systems*, Spriger-Verlag, Berlin (1992).
- [16] A. Karoui, *Wavelets: Properties and approximate solution of a second kind integral equation*, Comput. Math. Appl., **46** (2003) 263–277.
- [17] J.B. Keller, W.E. Olmstead, *Temperature of nonlinearly radiating semi-infinite solid*, Quart. Appl. Math., **29** (1972) 559–566.
- [18] X.Z. Liang, M.J. Che, *Solving second kind integral equations by Galerkin methods with continuous orthogonal wevelets*, J. Comput. Appl. Math., **136** (2001) 149–161.
- [19] K. Maleknejad, H. Derili, *Numerical solution of Hammerstein integral equation by using combination of spline-collocation method and Lagrange interpolation*, Appl. Math. Comput., **190** (2007) 1557–1562.
- [20] K. Maleknejad, K. Nouri, M. Nosrati Sahlan, *Convergence of approximate solution of nonlinear Fredholm-Hammerstein integral equations*, Commu. Nonlin. Sci. Numer. Simul., **15** (2010) 1432–1443.
- [21] K. Maleknejad, M. Tavassoli Kajani, *Solving second kind integral equations by Galerkin methods with hybrid Legendre and Block-Pulse functions*, Appl. Math. Comput., **145** (2003) 623–629.
- [22] J.S. Nadjafi, A. Ghorbani, *He's homotopy perturbation method: an effective tool for solving nonlinear integral and integro-differential equations*, Comput. Math. Appl., **58** (2009) 2379–2390.
- [23] W.E. Olmstead, R.A. Handelsman, *Diffusion in a semi-infinite region with nonlinear surface dissipation*, SIAM Rev., **18** (1996) 275–291.
- [24] G.M. Phillips, P.J. Taylor, *Theory and application of numerical analysis*, Academic Press, New York, (1973).
- [25] R. Piessens, *Computing integral transforms and solving integral equations using Chebyshev polynomial approximations*, J. Comput. Appl. Math., **121** (2000) 113–124.
- [26] L. Pretorius, D. Eyre, *Spline-Guss rules and the Nystrm method for solving integral equations quantum scattering*, J. Comput. Appl. Math., **18** (1987) 235–247.
- [27] R. Saadati, M. Dehghan, S.M. Vaezpour, M. Saravi, *The convergence of He's variational iteration method for solving integral equations*, Comput. Math. Appl., **58** (2009) 2167–2171.
- [28] H. Saeedi, M. Mohseni Moghadam, N. Mollahasani, G. N. Chuev, *A CAS Wavelet method for solving nonlinear Fredholm integro-differential equations of fractional order*, Commu. Nonlin. Sci. Numer. Simul., **16** (2011) 1154–1163.
- [29] G. Vainikko, A. Kivinukk, J. Lippus, *Fast solvers of integral equations of the second kind: wavelet methods*, J. Complex., **21** (2005) 243–273.

- [30] X.T. Wang, Y.M. Li, *Numerical solutions of integrodifferential systems by hybrid of general block-pulse functions and the second Chebyshev polynomials*, Appl. Math. Comput., **209** (2009) 266–272.
- [31] J.Y. Xiao, L.H. Wen, D. Zhang, *Solving second kind Fredholm integral equations by periodic wavelet Galerkin method*, Appl. Math. Comput., **175** (2006) 508–518.

Farshid Mirzaee

Department of Mathematics, Faculty of Science, Malayer University, Malayer, 65719-95863, Iran.

Email: f.mirzaee@malayeru.ac.ir

Elham Hadadiyan

Department of Mathematics, Faculty of Science, Malayer University, Malayer, 65719-95863, Iran.

Email: E.hadadiyan@gmail.com